

Sensitivity of Transit Searches to Habitable Planets

Andrew Gould, Joshua Pepper, D. L. DePoy

*Department of Astronomy, The Ohio State University, 140 W. 18th Ave., Columbus, OH
43210*

`gould,pepper,depoy@astronomy.ohio-state.edu`

ABSTRACT

Photon-limited transit surveys in V band are in principle about 20 times more sensitive to planets of fixed size in the habitable zone around M stars than G stars. In I band the ratio is about 400. The advantages that the habitable zone lies closer and that the stars are smaller (together with the numerical superiority of M stars) more than compensate for the reduced signal due to the lower luminosity of the later-type stars. That is, M stars can yield reliable transit detections at much fainter apparent magnitudes than G stars. However, to achieve this greater sensitivity, the later-type stars must be monitored to these correspondingly fainter magnitudes, which can engender several practical problems. We show that with modest modifications, the *Kepler* mission could extend its effective sensitivity from its current $M_V = 6$ to $M_V = 9$. This would not capture the whole M dwarf peak, but would roughly triple its sensitivity to Earth-like planets in the habitable zone. However, to take advantage of the huge bump in the sensitivity function at $M_V = 12$ would require major changes.

Subject headings: planets

1. Introduction

The one confirmed transiting planet discovered to date, HD209458b (Charbonneau et al. 2000), lies only $10 R_\odot$ from its host G star, and therefore well inside the so-called “habitable zone”, where water could exist in its liquid state. All ongoing transit surveys are primarily sensitive to such “hot Jupiters” because their large diameters give rise to relatively strong ($\sim 1\%$) signals, while their proximity to their host increases the probability of transits occurring (Howell et al. 2000; Brown & Charbonneau 1999; Mallen-Ornelas 2002; Udalski et al. 2002; Burke et al. 2002; Street et al. 2000, 2002).

While gas giants are not expected to themselves be habitable regardless of their location, they could have water-laden moons like Europa, Ganymede, and Callisto, which could support life if the planet lay in the habitable zone. Future ground-based surveys could be sensitive to such habitable-zone giant planets. Moreover, the planned *Kepler* mission¹ has as its primary goal the detection of planets down to Earth size in or near the habitable zone. The *Kepler* targets would be even more direct analogs of the Earth.

Here we investigate the sensitivity of transit surveys to planets in the habitable zone as a function of stellar type. We derive two remarkable conclusions. First, transit surveys are most sensitive to habitable planets orbiting middle M stars ($M_V \sim 12$), despite the fact that they are most sensitive overall to planets orbiting early G stars ($M_V \sim 4$). Second, to achieve this sensitivity to dim stars, the survey’s magnitude limit must be an extremely strong function of spectral type. Specifically, it must be about 7 magnitudes fainter for middle M stars than G stars. Since such variable requirements can significantly affect survey design, they deserve careful analysis and consideration. For example, we show that unless the *Kepler* mission extends its magnitude limit for M stars well beyond its current (type independent) limit of $V = 14$, it will lose most of its potential sensitivity to habitable planets.

2. Sensitivity to Habitable Planets

Pepper, Gould, & DePoy (2002) showed that if every star of luminosity L and radius R has a planet of radius r in a circular orbit of semi-major axis a , then a photon-noise limited transit survey will detect

$$N(L, R, r, a) = \frac{\Omega d_0^3}{3} n(L, R) \eta \frac{R_0}{a_0} \left(\frac{L}{L_0}\right)^{3/2} \left(\frac{R}{R_0}\right)^{-7/2} \left(\frac{a}{a_0}\right)^{-5/2} \left(\frac{r}{r_0}\right)^6 \quad (1)$$

planets. Here $n(L, R)$ is the number density of stars of the specified type, Ω is the angular area of the survey, and d_0 is the distance out to which an equatorial transit can just be reliably detected for the fiducial parameters L_0 , R_0 , r_0 , and a_0 . The numerical factor $\eta \simeq 0.719$ arises because the volume probed by the survey is smaller by a factor $y^{3/2}$ for non-equatorial transits, where y is the ratio of the transit chord to the stellar diameter. For the most part we will be comparing the detectability of planets of the same size around different stars. Hence, except when we are evaluating the normalization of equation (1) for a specific example, we will ignore the last factor.

¹<http://www.kepler.arc.nasa.gov/>

The equilibrium temperature of a planet (and thus whether or not water can exist in liquid phase on the planet surface) is the same if $a \propto L_{\text{bol}}^{1/2}$, where the bolometric luminosity L_{bol} is to be distinguished from L , the luminosity in the band of observation. Hence, the relative sensitivity of a transit survey to habitable planets is

$$N \propto nL^{3/2}R^{-7/2}L_{\text{bol}}^{-5/4}. \quad (2)$$

Here we assume that all planets have the same albedo and neglect atmospheric effects. If we compare, for example, G stars ($M_V = 5$) with middle M stars ($M_V = 12$), the ratios of the various factors are $N_{12}/N_5 \sim 6 \times 630^{-3/2} \times 4^{7/2} \times 80^{5/4} \sim 10$. Note that if we were comparing detectability at the *same semi-major axis* rather than the *same habitability*, the last factor would not have entered, and the ratio would have been 0.05. Hence, while the sensitivity to planets in general is completely dominated by G stars, the sensitivity to habitable planets is completely dominated by M stars. That is, the lower luminosities (and so smaller semi-major axes) combined with the smaller radii and greater numerical density of M stars more than compensate for the reduced photon counts.

In Figure 1, we show the sensitivity to habitable “Earths” and to Earths all at the same semi-major axis (taken to be 1 AU). That is, the histograms show the total number of planets N that will be detected as a function of M_V assuming that each star in the field has one Earth-size planet in the habitable zone or, respectively, one such planet at 1 AU. (The two histograms cross at $M_V = 5$ because the habitable zone is then at 1 AU). The absolute normalization of this plot is set according to the characteristics of the *Kepler* mission ($7.8 \times 10^8 \text{ e}^- \text{ hr}^{-1}$ at $V = 12$, $\Omega = 105 \text{ deg}^2$, $A_V = 0.3$, mission-total S/N=8 required for detection), but the form of the histogram would be the same for any photon-limited survey. The normalization for any other planet size should be multiplied by a factor $(r/r_{\oplus})^6$, and the normalization for any other fixed semi-major axis should be multiplied by $(a/\text{AU})^{-5/2}$ (see eq. [1]). The figure is constructed assuming that detection is in V band. The effect of substituting other bands is approximately to change the slope of the histogram. For example, since the slope of the main sequence is $dM_V/d(V - I) = 3.37$ (Reid 1991), substitution of I band would lead to an increase of slope $d\Delta \log N/dM_V = (3/2) \times 0.4/3.37 = 0.178$. That is, middle M stars would gain relative to G stars by an additional factor of $10^{7 \times 0.178} = 18$.

To compute these histograms, we follow the procedure of Pepper, Gould, & DePoy (2002). An important feature of the color-magnitude diagram is that while the main sequence is fairly narrow for $M_V > 6$, it broadens for brighter stars (due to faster stellar evolution), so that a star of a given M_V can have a large range of colors. Thus, for the well-defined lower main sequence, $M_V > 6$, we consider the luminosity function (Bessell & Stringfellow 1993; Zheng et al. 2001) in 1 mag bins, and evaluate the stellar radius at the center of each bin using the linear color-magnitude relation of Reid (1991), the color/surface-brightness

relation of van Belle (1999), and the *VIK* color-color relations from Bessell & Brett (1988).

On the other hand, for the upper main sequence $M_V < 6$ we evaluate the histograms directly using the Hipparcos catalog (ESA 1997). For example, the luminosity function for $M_V = 4$ is computed by summing $\sum_i [(4/3)\pi D_i^3]^{-1}$ over all stars within the Hipparcos completeness limit, $V < 7.3$, having $3.5 < M_V < 4.5$, and lying within 50 pc. The distance D_i is the minimum of 50 pc and the distance at which the star would have $V = 7.3$. Then, the constant-semi-major-axis histogram is computed by summing (and appropriately normalizing) $\sum_i L_i^{3/2} R_i^{-7/2} [(4/3)\pi D_i^3]^{-1}$, while the habitable-zone histogram is found by summing $\sum_i L_i^{3/2} R_i^{-7/2} L_{\text{bol},i}^{-5/4} [(4/3)\pi D_i^3]^{-1}$. The stellar radii are determined from Hipparcos/Tycho (B_T, V_T) photometry and the color/surface-brightness relation of Gould & Morgan (2003), ultimately derived from van Belle (1999). We evaluate L_{bol} using bolometric corrections as a function of $V - I$ color derived from Binney & Merrifield (1998) and Bessell & Brett (1988) at the bright end and Reid & Gilmore (1984) at the faint end.

3. Magnitude Limits for Habitable Planets

To achieve the sensitivities calculated in § 2, one must analyze the light curves of all the stars being probed. This statement would appear so obvious as not to be worth mentioning. However, as we now show, dim stars are being probed at substantially fainter apparent magnitudes than are their more luminous cousins. Hence, to avoid losing most of the sensitivity of a transit experiment one must set different magnitude limits for stars of different M_V .

Let m_{lim} be the apparent magnitude at which the survey achieves the minimum acceptable signal-to-noise ratio (S/N) for a planet of radius r and semi-major axis a circling an $M_V = 5$ star. Consider an identical planet circling an $M_V = 12$ star in the same orbit and at the same apparent magnitude. Since the star’s radius is a factor 4 smaller, the S/N will be a factor $4^2 \times 4^{-1/2} = 4^{3/2}$ times larger. The first factor comes from the fact that the planet occults a larger fraction of the stellar surface, the second from the reduced total time spent in transit (“duty cycle”) due to the smaller radius. Now, the bolometric luminosity of the dimmer star is down by a factor 80, so to keep in the habitable zone, the planet must be moved closer by a factor $80^{1/2}$. This increases the transit duty cycle by the same factor and so increases the S/N by $80^{1/4}$. Combining these two effects with the usual dependence of S/N on flux implies $\text{S/N} \propto \text{flux}^{1/2} L_{\text{bol}}^{-1/4} R^{-3/2}$. Hence, to maintain the same S/N, one must increase the magnitude limit by,

$$\Delta m_{\text{lim}} = 0.5 \Delta M_{\text{bol}} - 7.5 \Delta \log R. \quad (3)$$

For the example just given, $\Delta m_{\text{lim}} = 6.9$, i.e., a factor 570 in flux. Note that Δm_{lim} does not depend on the passband of observation. The dashed curve in Figure 1 shows the difference in the maximum distance modulus relative to the Sun at which a habitable planet (of given radius) can be detected. Specifically, $\Delta(m - M)_0 = (V_{\text{lim}} - M_V) - (V_{\text{lim},\odot} - M_{V,\odot})$. For V band observations, this offset is roughly constant over 10 magnitudes. That is, the limiting magnitude required to achieve the sensitivities shown by the bold histogram increases approximately in lock step with M_V . This behavior can have major consequences for the design of transit experiments.

4. Discussion

Photon-limited observations are not routinely achievable over very large dynamic ranges. A number of effects can intervene at fainter magnitudes. First, once the stars fall below the sky, the number of detected systems no longer falls as $L^{3/2}$ (see eq. [1]) but as L^3 . That is, once the sky is reached, detections would fall off by a factor ~ 4 per magnitude relative to the bold histogram in Figure 1. Second, if the exposure times are set so as not to saturate the brightest target stars (say 2 mag brighter than the mag limit appropriate for $M_V = 5$), then the flux levels will be a factor ~ 4000 times lower at the mag limit appropriate for $M_V = 12$. For many observing setups, this would render the flux levels comparable to the read noise. Third, a field that is very uncrowded at one magnitude may well be extremely crowded 7 magnitudes fainter, and this may interfere with doing photon-limited photometry. Hence, it is unlikely that all of the potential shown in Figure 1 can be achieved in any given practical experiment. Nevertheless, this potential is so great that it is worth thinking about how to achieve as much as possible.

We illustrate the type of trade-offs involved by considering the *Kepler* mission, the only transit experiment proposed to date whose primary target is planets in the habitable zone. The *Kepler* design calls for recording only stars $V < 14$, presumably in order to minimize data transmission. The mag limit required to detect Earth-like planets around Sun-like stars at S/N=8 is $V_{\text{lim},\odot} = 13.6$. Hence, from the dashed curve in Figure 1, the required mag limit at $M_V = 6$ is $V_{\text{lim}} = V_{\text{lim},\odot} - M_{V,\odot} + M_V + \Delta(m - M)_0 = 14.4$, i.e., already somewhat fainter than the $V = 14$ limit adopted by *Kepler*. The volume probed, and hence the detections, fall relative to the bold histogram by a factor $(10^{0.2})^3 \sim 4$ for each magnitude beyond $M_V = 6$. See Figure 2. Hence, assuming that every star has one planet in the habitable zone, only a total of 20 will be detected².

²Actually, we have not taken account of stellar variability in this analysis, nor of binaries. Each of these

Keeping the photometry information for stars to $V = 17$, for example, should increase the number of habitable-planet detections by almost a factor 3, while keeping stars with $V < 20$ would increase this number by a factor 9. See Figure 2. Of course, there are many obstacles to keeping information on stars this faint.

For example, there would be an enormous number of giant stars contaminating such a deep sample. In fact, however, distinguishing the later-type dwarfs from the much more numerous giants of similar colors is straightforward. As shown by Gould & Morgan (2003), they can easily be identified on a reduced proper motion (RPM) diagram constructed using data from USNO-A (Monet 1996, 1998) and 2MASS (Skrutskie et al. 1997). With USNO-B (Monet et al. 2002) it should be possible to extend coverage to fainter magnitudes and to achieve higher precision as well. The total number of such stars $M_V \leq 12$ is only $\sim 3 \times 10^4$, far fewer than the $\sim 10^5$ giants in the field that could be eliminated using the same RPM diagram (Gould & Morgan 2003).

A more difficult problem is crowding. The *Kepler* point spread function (PSF) is deliberately defocused to $10''$, meaning that the crowding limit is about 10 arcmin^{-2} . At the adopted *Kepler* sight line, $(l, b) = (69.6, 5.7)$, this limit is reached at $R_{\text{USNO}} \sim 18.2$ (as determined by a query of the USNO-A catalog). If *Kepler* were redirected to $(l, b) = (70, 15)$, the density of stars down to $R_{\text{USNO}} = 18.2$ would be 3.5 arcmin^{-2} , approximately 3 times lower. Hence, recovery of faint stars would be much easier. Presumably, *Kepler* has chosen to look right in the Galactic plane because of the higher overall star density. However, since the stars useful for transits mostly lie within 500 pc, a field at $b = 15^\circ$ would lie only 130 pc above the plane, where the density of such target stars is only slightly lower than at the plane. Hence, the crowding problem could be substantially ameliorated at small cost.

The last problem is sky. At high ecliptic latitude, the sky in space is $V \sim 23.3 \text{ arcsec}^{-2}$, or $V \sim 17$ in a $10''$ PSF. Hence, *Kepler* completeness appears to be fundamentally limited to stars $M_V < 9$. As shown in Figure 2, detections from stars at $M_V \geq 10$ would be highly suppressed. Only contracting the PSF (which has been deliberately defocused to improve the photometry) could overcome this difficulty. (Note that at $V \sim 17$, crowding would not be a serious problem even for the current Galactic-plane line of sight.)

While the above remarks apply specifically to *Kepler*, any transit experiment attempting to detect habitable planets would face similar constraints and trade-offs.

might plausibly reduce detections by 25%, the former by making transits unrecognizable, the later by making habitable planets dynamically unstable and by increasing the total light in the aperture. Hence, the true detection rate for this optimistic scenario should be reduced by roughly 50%.

Finally, we note that the results reported here are a strong function of planet size. As remarked in § 2, the absolute normalizations in Figure 1 scale as r^6 . By an argument similar to the one given in § 3, $\Delta m_{\text{lim}} = 10\Delta \log r$. For example, for $r = 0.63 r_{\oplus}$, the normalizations in Figure 1 would be reduced by a factor 16, while the required V_{lim} at each M_V would be brighter by 2 mag. Hence, in *Kepler*’s current configuration, it would retain full sensitivity to $M_V \sim 8$ stars, and so would be able to detect ~ 3 planets (see Fig. 2), while if its coverage were extended 3 mag to $V_{\text{lim}} \sim 17$, it could probe the M-dwarf peak, and so detect ~ 8 planets.

We thank Scott Gaudi for his detailed comments on the manuscript. This work was supported by grant AST 02-01266 from the NSF.

REFERENCES

- Bessell M.S., & Brett J.M. 1988, PASP, 100, 1134
- Bessell M.S., & Stringfellow G.S. 1993, ARA&A, 31, 433
- Binney, J., & Merrifield, M. 1998, Galactic Dynamics, (Princeton: Princeton Univ Press)
- Brown, T. M., & Charbonneau, D. 1999, BAAS, 31, 1534
- Burke, C.J., DePoy, D.L., Gaudi, B.S., Marshall, J.L., & Pogge, R.W. 2002, in ASP Conf. Ser., Scientific Frontiers in Research on Extrasolar Planets, eds. D. Deming and S. Seager, in press (astro-ph/0208233)
- Charbonneau, D., Brown, T. M., Latham, D. W., & Mayor, M. 2000, ApJ, 529, L45
- Gould, A., & Morgan, C.W. 2003, ApJ, 585, 000 (astro-ph/0209561)
- European Space Agency (ESA). 1997, The Hipparcos and Tycho Catalogues (SP-1200; Noordwijk: ESA)
- Howell, S. B., Everett, M., Davis, D. R., Weidenschilling, S. J., McGruder, C. H., III, Gelderman, R. 2000, BAAS, 32, 3203
- Mallen-Ornelas, G., Seager, S., Yee, H.K.C., Minniti, D., Gladders, M.D., Mallen-Fullerton, G., & Brown, T. 2002, ApJ, in press (astro-ph/0203218)
- Monet, D. G. 1996, American Astronomical Society Meeting, 188, 5404.

- Monet, D. G. 1998, American Astronomical Society Meeting, 193, 112003
- Monet, D. G., et al. 2002, AJ, in press (astro-ph/0210694)
- Pepper, J., Gould, A., & DePoy, D.L. 2002, ApJ, submitted (astro-ph/0208042)
- Reid, N. 1991, AJ, 102, 1428
- Reid, I.N., & Gilmore, G.F. 1984, MNRAS, 206, 19
- Skrutskie, M.F. et al. 1997, in The Impact of Large-Scale Near-IR Sky Survey, ed. F. Garzon et al. (Kluwer: Dordrecht), p. 187
- Street, R. A. et al. 2000, in ASP Conf. Ser., Vol. 219, Euroconference on Disks, Planetesimals and Planets, eds. F. Garzon, C. Eiroa, D. de Winter, & T. J. Mahoney (San Francisco: ASP), 572
- Street, R. A. et al. 2002, in ASP Conf. Ser., Scientific Frontiers in Research on Extrasolar Planets, eds. D. Deming and S. Seager, in press (astro-ph/0208233)
- Udalski, A. et al. 2002, AcA, 52, 1
- van Belle, G.T. 1999, PASP, 111, 1515
- Zheng, Z., Flynn, C., Gould, A., Bahcall, J.N., Salim, S. 2001, ApJ, 555, 393




Fig. 1.— Total number of planets detected in the habitable zone (*bold histogram*) and in 1 AU orbits (*solid histogram*) as functions of absolute magnitude M_V (assuming every star has one such planet). The absolute normalizations have been set to Earth-size planets and to the characteristics of the *Kepler* mission, but the form is completely general assuming observations in V band. In I band, the slope would be tilted upward toward faint stars. Extinction has been taken into account, but stellar variability and binarity have not. The calculation uses different methods for early ($M_V \leq 6$) and late ($M_V \geq 6$) stars. The solid and dashed histograms compare the two methods in the one bin of overlap. The *dashed curve* shows the difference (relative to Sun-type stars) in the limiting distance modulus to which stars of different M_V must be monitored to achieve the sensitivity shown by the “habitable zone” histogram. Since this curve is basically flat, dimmer stars must be observed to the same distance and hence to much fainter apparent magnitudes.




Fig. 2.— Total number of planets detected normalized to characteristics of *Kepler* and under various assumptions. *Bold solid histogram*: each star has one Earth-size planet in the habitable zone and all stars are monitored regardless of magnitude (same as Fig. 1). *Bold dashed histogram*: same planet distribution, but now only stars $V < 14$ are monitored. *Thin dashed histogram*: same planet distribution, $V < 17$ stars are monitored. *Dotted histogram*: same planet distribution, $V < 20$ stars are monitored. *Thin solid histogram*: planets have radius $r = 0.63 r_{\oplus}$ and only $V < 14$ stars are monitored.